



CGCTCAGGATATGAGTCTCCGCTAGTTCATCGGATCCTGGGCTGATATATATAGCTCGATCGATC  
 TTCTCTATATGCGGATTCGCTATATATACACACACACACACACACACACACACACACACACACAC  
 ACCGATCTAT  
 CACAGACATACGCTAT

Nucleotide

PubMed	Nucleotide	Protein	Genome	Structure	PMC	Taxonomy	OMI
Search <input type="text" value="Nucleotide"/> for <input type="text"/>							
Limits		Preview/Index		History		Clipboard	
Display	<input type="text" value="default"/>	Show:	<input type="text" value="20"/>	Send to	<input type="text" value="File"/>	Get Subsequence	

☐ 1: U65943. Chlamydomophila abo...[gi:1617508]

Links

LOCUS CPU65943 6234 bp DNA linear BCT 07-  
 DEFINITION Chlamydomophila abortus clone S21 POMP91B precursor and POMP9  
 precursor, genes, complete cds.  
 ACCESSION U65943  
 VERSION U65943.1 GI:1617508  
 KEYWORDS .  
 SOURCE Chlamydomophila abortus  
 ORGANISM Chlamydomophila abortus  
 Bacteria; Chlamydiae; Chlamydiales; Chlamydiaceae; Chlamydo  
 REFERENCE 1 (sites)  
 AUTHORS Longbottom,D., Russell,M., Jones,G.E., Lainson,F.A. and  
 Herring,A.J.  
 TITLE Identification of a multigene family coding for the 90 kDa  
 of the ovine abortion subtype of Chlamydia psittaci  
 JOURNAL FEMS Microbiol. Lett. 142 (2-3), 277-281 (1996)  
 MEDLINE 96406378  
 PUBMED 8810511  
 REFERENCE 2 (sites)  
 AUTHORS Longbottom,D., Russell,M., Dunbar,S.M., Jones,G.E. and Herr  
 TITLE Molecular cloning and characterization of the genes coding  
 highly immunogenic cluster of 90-kilodalton envelope protei  
 the Chlamydia psittaci subtype that causes abortion in shee  
 JOURNAL Infect. Immun. 66 (4), 1317-1324 (1998)  
 MEDLINE 98187897  
 PUBMED 9529048  
 REFERENCE 3 (bases 1 to 6234)  
 AUTHORS Longbottom,D.  
 TITLE Direct Submission  
 JOURNAL Submitted (02-AUG-1996) Moredun Research Institute, Interna  
 Research Centre, Pentlands Science Park, Bush Loan, Penicu  
 Midlothian EH26 0PZ, UK  
 FEATURES Location/Qualifiers  
 source 1..6234  
 /organism="Chlamydomophila abortus"  
 /mol\_type="genomic DNA"  
 /strain="S26/3"  
 /specific\_host="Ovis aries"  
 /db\_xref="taxon:83555"  
 /clone="S21 and S432"

/clone\_lib="lambda EMBL-3"  
/dev\_stage="elementary body"  
/note="submitted as Chlamydia psittaci"  
-35 signal 543..548  
/note="putative"  
/citation=[2]  
-10 signal 575..580  
/note="putative"  
/citation=[2]  
RBS 608..614  
/note="putative"  
/citation=[2]  
CDS 619..3159  
/note="encodes putative 90,847 Da outer membrane p  
/citation=[2]  
/codon\_start=1  
/transl\_table=11  
/product="POMP91B precursor"  
/protein\_id="AAC15923.1"  
/db\_xref="GI:1617509"  
/translation="MKHPVYWFLISSSLLASNSLSFAQVTNETLTSSDSY  
FEVKETTSGAIYTCGNVCISYAGKDSPLNKSCFSETTENLSFIGNGYTL  
SSHPGAISVSGTNKTLDISGFSLFSCAYCCPPGTTGYGAIQTKGTTTLKD  
NCSTAEGGAIQCKSSSSTAELKLENNKNLVFSENSSKEKGGAAYADKLT  
SNNSVSHNSSPKGGAICIKSDSGECSLTANLGDIITFDGNKIITNGGSPT  
GSGGKFTKLNAKEGFGIFFYDPIANTGGSTEIELNKTESDTTYTGKIVFS  
KTVPANLKSIFYKQPLKIGAGSLVLKDGVTLEAKKITQTKGSTVMDLGT  
ETITLTNLDINIASLGGGGGTAPAKLATNTASQAISIAAVNLVNTDSNTY  
KSFSAITATTSSSTVTPPETNLKNYTPPTHYGYQGNWTVTWKQGSSAQEK  
TGYSPPNPERVGS LVPNTLWGAFSDTRAIQNLMDISVNGADYSRGFWVSSL  
SDTKRKFRHHSAGYALGVYAQTPSDDVCSAAFCQLFGKDKDYFVSKNSST  
QHISYWNTWNTLLQNTLGAEAPLVLNAQLTYCHASNMMKTNMTNTYTPKN  
DWGNDCFGVEFGAKAPIETASLLFDMYSPFVKLQLVHAHQDDFKENNSDQ  
LTNLSMPIGVKLEKFSHKDTASYNLTLAYAPDIVRSNPDCTASLLVSPTS  
NLARHAFILQAGNYLALTRNTELFSGFELRGSCRTYNIDLGSKIQF"  
sig peptide 619..666  
/note="putative"  
mat peptide 667..3156  
/product="POMP91B"  
/note="putative"  
misc feature 748..1202  
/note="identical to clone 99 antigen encoded by Ge  
Accession Number U59307"  
/citation=[1]  
/citation=[2]  
/evidence=experimental  
stem loop 3173..3198  
/citation=[2]  
/function="putative 11-base dyad terminator"  
-35 signal 3222..3227  
/note="putative"  
/citation=[2]  
-10 signal 3246..3251  
/note="putative"

/citation=[2]  
3279..3285  
/note="putative"  
/citation=[2]  
3290..5809  
/note="encodes putative 89,836 Da outer membrane p  
/citation=[2]  
/codon\_start=1  
/transl\_table=11  
/product="POMP90B precursor"  
/protein\_id="AAC15924.1"  
/db\_xref="GI:1617510"  
/translation="MKHPVYWFLISSSLFASNSLSFANDAQTALTPSDSY  
FQVKETSSGTTYTCEGNVCISFAGKDSGLKKSCFSATDNLTFLGNGYTL  
SNPGAINVQGGKTLGISGFSLFSCAYCPPGTTGYGAIQTKGNTTLKDNS  
STAEGGAIQCKGSSDAELKIENNQNLVFSSENSSTSKGGAIYADKLTIVSG  
SVSNGSSPKGGAISIKDSSGECSLTADLGDITFDGNKI IKTSGGSSTVTR  
KFTKLRAKDGFGIFFYDPITGGGSDELNINKKETVDYTGKIVFSGEKLS  
LASTFNQPI TLSAGSLVLKDGVSVTAKQVTQEAGSTVMDLGTTLQTPSS  
NLDINIASLGGGGGTSPAKLATNTASQAITINAVNLVDADGNAYEDPILA  
VATTNASTVTQPTDNL TNYVPPTHYGYQGNWTVTWD TETATKTATLTWEQ  
RQGPLVPNTLWGAFSDLRAIQNLMDISVNGADYHRGFVWSGLANFLHKSG  
HNSAGYALGVYAKTPSDDIFSA AFCQLFGKDKDYLVS KNNANIYAGSLYY  
WQNLLQNTIGAEAPLVLNAQLTYCHASN DMKTNMTTTYAPRKTTYAEIKG  
VELGATVP IQTESSLLFDMYSPFLKFQLVH THQDDFKENNSDQGRYFESS  
IGIKFERFANNDTASYHVTAAYSPDIVRSNP DCTTSLLVSPDSAVVWTKA  
MLQAGNYLSLSH NIEIFSQGFELRGSSRTYNVDL GSKIQF"  
sig peptide 3290..3337  
/note="putative"  
mat peptide 3338..5806  
/product="POMP90B"  
/note="putative"  
misc feature 4220..4435  
/note="identical to clone 31 antigen encoded by Ge  
Accession Number U59306"  
/citation=[1]  
/citation=[2]  
/evidence=experimental  
stem loop 5823..5848  
/citation=[2]  
/function="putative 11-base dyad terminator"  
stem loop 5910..5938  
/note="second stem loop structure"  
/citation=[2]

BASE COUNT 1927 a 1429 c 1152 g 1726 t

ORIGIN

```
1 ctgcagggttc agtttttttta taacttcaac atcttctttt gaaaaaacat aatcgg
61 tttagatagtg aaaacacaat gagataaagt ttttagttca tcaacggtta gctgag
121 aaacttttcc agtttctgtc tactttgttg attaccatgt atcacaaaat aagcgc
181 gctttccaaa cagtagggat caagattcca aaaatcctca tattgatctt gcggga
241 cgcaccttca caaaatatct ttgagccaaa tccttctaca gcaataacaa acacaa
301 agcaaacagc gaccttggca tcataaatgg tacttcctga gtatctttta tctgaa
361 ctgatgtgag aattatacca cggttcctct catattgtga ttctttatth actcat
421 tagatcacag atagatcatg aaaagaaata aggaatagtt tctatttcat catctt
```


481	aagcctacct	atctttgcaa	tcacaaatgc	gtatatgtaa	aaggttggag	ttgtgt
541	attagagacg	atccccttga	ctttttaata	tttttatgat	cgacttggga	tatctc
601	cacccaaagc	atgaatttat	gaaacatcca	gtctactggt	tcttaatatc	ctcaag
661	ctcgcctcga	attcttttgag	ctttgctcag	gtaacaaatg	aaaccttaac	ttcctc
721	agctataatg	gaaatgtcac	ctctgatgaa	ttcgaggtaa	aagaaactac	atcagg
781	atttatactt	gcgaaggtaa	tgtgtgcac	tcctatgcag	ggaaagattc	tcctct
841	aaaagttggt	tctcagaaac	tactgaaaat	ctttctttca	taggaaatgg	ttacac
901	tgttttgata	atattactac	acaatctagt	caccccgag	ctattagtgt	tagtgg
961	aataaaacct	tagacatctc	aggattttcc	ttattttcat	gtgcctattg	ctgcc
1021	ggaaccaccg	gttacggagc	tatacagact	aaaggcacca	caactttaaa	agataa
1081	agtcttgtct	tccataaaaa	ctgttcgaca	gcagagggtg	gcgctattca	gtgtaa
1141	agcagctcta	ctgctgagtt	aaagctagaa	aataataaaa	atcttgtttt	ctcaga
1201	tcttctaagg	aaaaaggcgg	ggctatttat	gccgataaac	tcaccattgt	ctcagg
1261	ccgacgttat	tttctaacaa	ctccgtatcc	cacaattcat	cccctaaagg	cggggc
1321	tgtatcaaag	attcagatgg	tgaatgtagt	ctaacggcga	atcttgggga	tattac
1381	gatggaaaca	aaatcatcac	aactaatggt	ggaagtccta	cagtaacaag	aaattc
1441	gatctcggct	ctggtggaaa	attcacaaaa	ctcaatgcta	aagaaggttt	tgggat
1501	ttttatgacc	ctatcgctaa	cacaggagga	tctacagaaa	tcgagctaaa	taaaac
1561	agcgacacta	cttatacagg	aaagatcgtc	ttctcaggag	aaaagttatc	tgatga
1621	aaaacggttc	cggccaatct	gaaatcatat	ttcaaacaac	ccctaaaaat	cgggtgc
1681	tcttttagtcc	ttaaagatgg	tgtcactcta	gaagcaaaaa	aaattacaca	gacaaa
1741	tctaccgttg	tcattggatct	agggactacg	ttacaaacac	cgtcctccag	tggaga
1801	attaccttaa	cgaatttgga	tattaacatc	gcctcgttgg	gggggggggg	gggtac
1861	ccagccaaac	tcgcaacgaa	tacagcaagc	caagctataa	gtattgccgc	tgttaa
1921	gtcaataactg	acagcaatac	ttatgaagat	cccatactct	ccgcatcaaa	atcttt
1981	gcaataacag	ccacaacctc	ctcaagcaca	gtcacacctc	ccgaaaccaa	tctaaa
2041	tatactcctc	ctacccatta	cgtttatcaa	ggtaattgga	cggtcacttg	gaaaca
2101	tcaagcgccc	aagaaaaaac	agccactcta	acttggggaa	aaacgggtta	ctctcc
2161	ccagaacgtg	taggatcttt	agtgcctaac	actctctggg	gcgcattctc	tgatac
2221	gctatacaaa	acttaattgga	tatcagcgtc	aatggcgctg	actactcaag	aggttt
2281	gtatctagtc	tagccaactt	cctaaacaaa	agcggttctg	acactaaacg	caagtt
2341	caccatagtg	ctgggtatgc	tttaggcgct	tatgcacaaa	ctccttctga	tgacgt
2401	agcgcagctt	tctgccaact	cttcggaaaa	gacaaagact	atttcgtctc	caaaaa
2461	tccactatct	acgcaggatc	tatctattat	caacatattt	cttattggaa	tacctg
2521	acgttgctac	aaaatactct	cgggtgcggag	gccccgttag	tcctgaatgc	gcagtt
2581	tattgtcatg	cttcaaataa	catgaaaacc	aacatgacaa	atacatatac	tcctaa
2641	gtcacaccct	cagaaatcaa	agggtgattgg	ggaaatgatt	gtttcggagt	cgagtt
2701	gctaaagcgc	ccatagaaac	ggcatctctc	ctcttcgata	tgtattcacc	tttcgt
2761	ctacaacttg	tgcatgcgca	ccaagatgac	tttaaggaaa	acaatagcga	tcaagg
2821	tactttgaaa	gtaataacct	caccaactta	tctatgccca	taggtgtgaa	attgga
2881	ttctctcaca	aggataccgc	ttcttacaa	cttaccctgg	cttatgctcc	agatat
2941	agaagcaacc	ctgactgtac	tgctctctg	ttagtgaacc	caacctctgc	tgtctg
3001	actaaagcta	ataaccttgc	gcggcatgct	ttcatattac	aagcaggaaa	ttactt
3061	ttaactcgca	acacggagct	cttcagtcac	ttcggtttcg	aactccgagg	ttcttg
3121	acctataaca	tagatctcgg	atcgaagatc	cagtttcta	ctcatccacc	ttccct
3181	cgtagacacg	gagcagggcc	tccctttgtc	tactctagct	ggtgaccttg	actttt
3241	atthtttatga	tcgagttagg	atatctcaaa	tcacccaaag	catgaattta	tgaac
3301	agtctactgg	ttcttaatat	cctcgagcct	atthgcctcg	aattctttga	gcttcg
3361	cgacgctcaa	acagccttaa	ctccctccga	tagctataat	ggaaatgtga	cctctg
3421	gttccaggta	aaagaaactt	catcaggaac	aacgtatact	tgtgaaggca	atgtgt
3481	ctcctttgca	gggaaagatt	caggtctaaa	gaaaagttgt	ttctcagcta	ctgata
3541	taccttccta	ggaaacgggt	atactctttg	ctttgataat	attactacta	cagcta
3601	ccccggagcc	attaatgttc	aagggtcaagg	aaaaacctta	ggcatctcag	gatttt
3661	atthttcatgt	gcttattgtc	ctccaggcac	aactggttac	ggagctatac	agacta

3721 caacacaact ttaaaagata actctagtct tgtcttccat aaaaactgct caacag  
3781 aggtggggct atccagtgt aaggaagcag tgatgctgaa ttaaaaatag aaaata  
3841 gaatctggtt ttctcagaaa actcctccac ttcaaaaggc ggggctattt atgctg  
3901 actcaccatt gtctcaggtg ggcctacatt attttctaac aactctgtat ccaacg  
3961 atcccctaaa ggcggagcta ttagcataaa agattcaagt ggtgaatgta gcctaa  
4021 tgatctcgga gatattacct tcgatgggaa caaaatcatc aaaactagtg gtggaa  
4081 tacagtaaca agaaattcca tagatctcgg cacagggaaa ttacaaagc tacgtg  
4141 agacggcttc ggaattttct tctatgacct tattactggg ggaggatctg atgaac  
4201 cattaataaa aaagaaactg ttgattatac aggaaagatc gtcttctcag gtgaaa  
4261 atccgatgaa gaaaaagcac gagcggaaaa cctagcttct actttcaacc aaccga  
4321 attatcagca ggatctcttg tacttaaaaga tgggtgatct gtaaccgcaa aacaag  
4381 gcaggaagcg ggatctaccg ttgtcatgga tctagggacc acattacaga cgcctt  
4441 aggtggagaa accatcaccc taactaatct agatattaac atcgctctgt tggggg  
4501 ggggggtacc tctcctgcta aactcgcaac aaatacagca agtcaagcta taacta  
4561 cgctgtcaat ctagtcgatg ctgatggcaa tgcttatgaa gatcctattc ttgcta  
4621 taaacctttc acagcaatag tagctacaac taacgctagt acagtcacac agccta  
4681 taatctaaca aattatgtcc ctctactca ttacggttac caaggaaatt ggacag  
4741 ttgggacacc gaaacagcta caaaaacagc cactctaact tgggaacaaa ctggct  
4801 ccctaaccga gaacgtcaag gacctttagt cccgaatact ctttgggggt cattct  
4861 cctcagagct atacaaaact taatggatat tagcgtcaat ggcgctgact accata  
4921 tttttgggta tccggtctag ctaacttctt acacaaaagt ggctctgata ctaaac  
4981 gttccgtcac aatagcgccg gatacgcttt aggcgtctac gcaaaaactc cttctg  
5041 tattttcagt gcggctttct gccaaactct cggaaaggac aaagactatt tagtgt  
5101 aaacaacgcc aacattttac caggttctct ctattatcag catatctcct attgga  
5161 ttggcagaat ctgctacaaa acactatcgg tgcagaagct ccggtagtcc ttaacg  
5221 gttaacttat tgatcatgct caaacgacat gaaaaccaac atgacgacta cttacg  
5281 tcgtaaaaca acgtatgcag aaatcaaggg tgattggggg aacgattgtt tcggag  
5341 gcttgggtgca actgtgccta tccaaacaga atcttctctc ctattcgata tgtact  
5401 tttcctgaag tttcaacttg tgcatacgca ccaagatgac ttaaggaaa acaata  
5461 tcagggaaga tacttcgaaa gcagcaatct caccaacctt tctctgccta tcggca  
5521 gtttgagaga tttgctaaca acgatacagc ttcttatcat gtcactgctg cttatt  
5581 tgatatacgt agaagtaacc ctgactgtac tacttctctg ttagtaagcc ccgact  
5641 tgtctgggta acgaaagcca acaaccttgc gcgaagcgcc ttcagctac aagcag  
5701 ctacttgtct ttaagtcaca acatagaaat cttcagccag ttcggtttcg agctca  
5761 atcttcacga acctataacg tagatctcgg atcgaagatc cagttctaata ctcac  
5821 tctcctacct cgttcccacg gggtagggcc tcctcggcct gtgacgctaa gactct  
5881 caaacatgaa aaagctcaca atcacgaaga atttctaaga gtgtacttct tggaaa  
5941 ccgctaactt ttttaaacag tatgatatct ttggacatcg ttaactctat agaagt  
6001 gctttaatat cacctttaag taaagctttc gataatagtg taactacctt ctgctg  
6061 aaccgtttta aaggtcgtgc tccaaaagca ctatcatacc cctgctcact gagata  
6121 ataacagaat catcccatgt taacgtgact cgtctttcta acatgcgttg ggcaac  
6181 cgcactctgaa tgccgacgat ttttacaata tcttcggtac ttaaaggaat aaaa

//

[Disclaimer](#) | [Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)

Mar 17 2003 10:55:57



PubMed	Nucleotide	Protein	Genome	Structure	PMC	Taxonomy	OMI
Search <input type="text" value="Nucleotide"/> for <input type="text"/>							
Limits		Preview/Index		History		Clipboard	
Display	<input type="text" value="default"/>	Show:	<input type="text" value="20"/>	Send to	<input type="text" value="File"/>	Get Subsequence	

# ☐ 1: U65942. Chlamydomophila abo...[gi:1617505]

[Links](#)

LOCUS CPU65942 6110 bp DNA linear BCT 07-  
 DEFINITION Chlamydomophila abortus strain S26/3 POMP91A and POMP90A prec  
 genes, complete cds.  
 ACCESSION U65942  
 VERSION U65942.1 GI:1617505  
 KEYWORDS .  
 SOURCE Chlamydomophila abortus  
 ORGANISM Chlamydomophila abortus  
 Bacteria; Chlamydiae; Chlamydiales; Chlamydiaceae; Chlamydo  
 REFERENCE 1 (sites)  
 AUTHORS Longbottom,D., Russell,M., Jones,G.E., Lainson,F.A. and  
 Herring,A.J.  
 TITLE Identification of a multigene family coding for the 90 kDa  
 of the ovine abortion subtype of Chlamydia psittaci  
 JOURNAL FEMS Microbiol. Lett. 142 (2-3), 277-281 (1996)  
 MEDLINE 96406378  
 PUBMED 8810511  
 REFERENCE 2 (sites)  
 AUTHORS Longbottom,D., Russell,M., Dunbar,S.M., Jones,G.E. and Herr  
 TITLE Molecular cloning and characterization of the genes coding  
 highly immunogenic cluster of 90-kilodalton envelope protei  
 the Chlamydia psittaci subtype that causes abortion in shee  
 JOURNAL Infect. Immun. 66 (4), 1317-1324 (1998)  
 MEDLINE 98187897  
 PUBMED 9529048  
 REFERENCE 3 (bases 1 to 6110)  
 AUTHORS Longbottom,D.  
 TITLE Direct Submission  
 JOURNAL Submitted (02-AUG-1996) Moredun Research Institute, Interna  
 Research Centre, Pentlands Science Park, Bush Loan, Penicui  
 Midlothian EH26 0PZ, UK  
 FEATURES Location/Qualifiers  
 source 1..6110  
 /organism="Chlamydomophila abortus"  
 /mol\_type="genomic DNA"  
 /strain="S26/3"  
 /specific\_host="Ovis aries"  
 /db\_xref="taxon:83555"  
 /clone="S112"

/clone\_lib="lambda EMBL-3"  
/dev\_stage="elementary body"  
/note="submitted as Chlamydia psittaci"  
-35 signal 3..8  
/note="putative"  
/citation=[2]  
-10 signal 33..38  
/note="putative"  
/citation=[2]  
RBS 66..72  
/note="putative"  
/citation=[2]  
CDS 77..2620  
/note="encodes putative 90,707 Da outer membrane p  
/citation=[2]  
/codon\_start=1  
/transl\_table=11  
/product="POMP91A"  
/protein\_id="AAC15921.1"  
/db\_xref="GI:1617506"  
/translation="MKHPVYWFLISSSLIVSNSLYSEEPDQKTLTSAHSY  
FNPLSTSNSNGTIYTCTGNICIAIYAGLDGSGLSSSCFTDTAGNLSFLGNG  
TTQSSHPGAISVSGTNKTLDISGFSLFSCAYCPPGATGYGAIKAVGNTTI  
HKNCSTGEGGAIQCKASSSEAELEKIENNQNLFVFAENSSSSSSGGAIYADKL  
LFSNNSVSASSPKGGAICIKDSGGECSLTADLGDITFDGNKIIKTNGGSP  
LGSSGKFTKLNAKEGFGIFFYDPITGGGSDELNINKQDVTVDYTGKIVFSG  
KVAANLKSDFKQPLKIGSGSLILKDGVTLETKSFTQTEGATVMDLGTTL  
TITLTNLDINVASLGGGGVAPDPAKVEATTESKTVTINAVNLVDDNGNAY  
QPFTAIEVRSGSSGSITKPTTNLENYTPPTHYGYQGNWTVTWKQGSSAQE  
QTGYSPNPERQGS LVPNTLWGSFSDIRAIQNLMDISVNGADYHRGFWVSG  
GSDTKRKFRHNSAGYALGVYAQTPSEDVFSA AFCQLFGKDKDYLVSKNSS  
YQHISYWNTWNTLLQNTLGAEAPLV LNAQLAYCHASNMMKTNMTDTYAPP  
GDWGNDCFGVEFGAKAPIETASLLFDMYSPFVKLQLVHAHQDDFKENNSD  
NLTNLSMPIGVKLEKFSHKDTASYNLT LAYAPDIVRSNPDCTASLLVSPT  
NNLARHAFILQAGNYLALTRNTELF SQFGFELRGSCRTYNIDLGSKIQF"  
stem loop 2634..2659  
/citation=[2]  
/function="putative 11-base dyad terminator"  
-35 signal 2683..2688  
/note="putative"  
/citation=[2]  
-10 signal 2707..2712  
/note="putative"  
/citation=[2]  
RBS 2740..2746  
/note="putative"  
/citation=[2]  
CDS 2751..5270  
/note="encodes putative 89,836 Da outer membrane p  
/citation=[2]  
/codon\_start=1  
/transl\_table=11  
/product="POMP90A precursor"  
/protein\_id="AAC15922.1"

/db\_xref="GI:1617507"  
/translation="MKHPVYWFLISSSLFASNSLSFANDAQTALTPSDSY  
FQVKETSSGTTYTCEGNVCISFAGKDSGLKKSCFSATDNLTLGNGYTL  
SNPGAINVQGGKTLGISGFSLFSCAYCPPGTTGYGAIQTKGNTTLKDNS  
STAEGGAIQCKGSSDAELKIENNQNLFSENSSTSKGGAIYADKLTIVSG  
SVSNGSSPKGGAISIKDSSGECSLTADLGDITFDGNKI IKTSGGSSTVTR  
KFTKLRAKDGFGIFFYDPITGGGSDELNINKKETVDYTGKIVFSGEKLS  
LASTFNQPI TLSAGSLVLKDGVSVTAKQVTQEAGSTVMDLGTTLQTPSS  
NLDINIASLGGGGGTSPAKLATNTASQAITINAVNLVDADGNAYEDPILA  
VATTNASTVTQPTDNL TNYPPTHYGYQGNWTVTWD TETATKTATLTWEQ  
RQGPLVPNTLWGA FSDLRAIQNLMDISVNGADYHRGFVWSGLANFLHKS  
G HNSAGYALGVYAKTPSDDIFSA AFCQLFGKDKDYLVS KNNANIYAGSLY  
Y WQNL LQNTIGAEAPLV LNAQLTYCHASN DMKTNMTT TYAPRKTTYAEIK  
G VELGATVPIQTESSLLFDMYSPFLKFQLVH THQDDFKENNSDQGRYFESS  
I GIKFERFANNDTASYHVTAAYS PDIVRSNP DCTTSLLVSPDSAVWVTKA  
MLQAGNYLSLSH NIEIFSQFGFELRGSSRTYNVDL GSKIQF"

sig peptide

2751..2798

/note="putative"

mat peptide

2799..5264

/product="POMP90A"

/note="putative"

misc feature

3681..3896

/note="identical to clone 31 antigen encoded by Ge  
Accession Number U59306"

/citation=[1]

/citation=[2]

/evidence=experimental

stem loop

5284..5309

/citation=[2]

/function="putative 11-base dyad terminator"

BASE COUNT        1856 a    1415 c    1163 g    1676 t  
ORIGIN

1	tctagacaat	ccccttgact	tttaattatt	tttatgatcg	gtttggggata	tctcaa
61	cccaaagcat	gaatttatga	aacatccagt	ttactggttt	ttaatatcct	cgagct
121	cgtctcgaac	tcattatact	ccgaggaacc	agatcaaaaa	accttgacct	ctgcgc
181	ctacaatgga	aacacgaata	gtgagccatt	caatccatta	tcaactagca	attcaa
241	tacaatatac	acgtgtacag	ggaatatatg	tattgcctat	gcgggggttag	atggct
301	cttatccagt	agttgtttca	ctgatactgc	tggaaacctt	tccttcttgg	gaaatg
361	cacactgtgt	tttgataata	ttactacaca	atctagtcac	cccggagcta	ttagtg
421	tggtagcaat	aaaaccttag	acatctcagg	attttcctta	ttttcatgtg	cctatt
481	tccaggagca	accggctacg	gagccatcaa	agctgtaggg	aataccacta	tcaaag
541	ctctagtctc	gtcttccata	aaaactgctc	tacaggagaa	ggtggggcta	ttcagt
601	agcaagcagt	agtgaagctg	aattaaaaat	agaaaataat	caaaatcttg	ttttcg
661	aaactcctcc	tcttcaagcg	gcggggctat	ttatgctgat	aaactcacca	ttgtct
721	tggacctacg	ttattttcta	ataactccgt	atccgcttct	tcacctaaag	gtggag
781	ttgcataaaa	gattcaggtg	gtgaatgtag	cttaaccgct	gatctcggag	atatca
841	tgatgggaac	aaaatcatca	aaactaatgg	tggaaagtcct	acagtaacaa	gaaatt
901	cgatctcggc	tctagcggaa	aatttacaaa	actaaatgct	aaagaaggtt	tcggga
961	cttctatgac	cctattactg	gaggaggatc	tgatgaatta	aatattaata	aacaag
1021	tgttgattat	acaggcaaga	tcgtcttctc	tggtgaaaga	ttatcagatg	aagaaa
1081	ggttgcgggc	aatctgaaat	cagatttcaa	acaaccctta	aaaatcggtt	ccggat
1141	aatccttaaa	gatggtgtca	ctctagaaac	gaaatccttc	acgcaaacag	agggcg
1201	cgttgtcatg	gacctaggga	ctacgttaca	aacgccttcc	tcaggtggag	aaacca
1261	cctaactaat	ttggatatta	acgtcgcctc	attgggaggg	gggggggtcg	ctccag




1321	tgctaaagtc	gaagcaacaa	ctgaaagtaa	aacagtcaca	atcaatgctg	tcaatc
1381	cgatgacaat	ggcaatgcct	acgagtatcc	tatacttgct	gcattctcagc	ctttca
1441	aatagaagta	agatctgggt	ctagcggctc	tattactaaa	cccacaacca	atctag
1501	ctatacccct	cctacccatt	acggttatca	aggtaattgg	acggtcactt	ggaaac
1561	atcaagcgcc	caagaaaaaa	cagccactct	aacttgggaa	caaactgggt	actccc
1621	cccagaacgt	caaggatctt	tagttccgaa	tactctctgg	ggttccttct	ctgaca
1681	agctatacaa	aacttaaatg	atatcagcgt	caatgggtgct	gattaccata	gagggt
1741	ggtatccgg	ctaggaaact	tcttacataa	aagtggctca	gatactaaac	gaaaat
1801	ccacaatagt	gccggatacg	ccttaggcgt	ctacgcgcaa	actccttcag	aagacg
1861	cagtgcggct	ttctgccaac	tcttcggaaa	agacaaaagac	tatttagtct	ctaaaa
1921	ttccactgtc	tatgcagggt	ccatctatta	tcaacatatt	tcttattgga	atacct
1981	tacgttgcta	caaaatactc	tcggtgcgga	agctccgtta	gtccttaacg	cacagt
2041	ttattgtcat	gcttctaata	atatgaagac	aaacatgaca	gacacctacg	ctcctc
2101	aacaacatat	tcagaaatca	aaggtgattg	gggaaatgat	tgtttcggag	tcgagt
2161	agctaaagcg	cccatagaaa	cggcatctct	cctcttcgat	atgtattcac	ctttcg
2221	gctacaactt	gtgcatgcgc	accaagatga	ctttaaggaa	aacaatagcg	atcaag
2281	atactttgaa	agtaataacc	tcaccaactt	atctatgccc	ataggtgtga	aattgg
2341	attctctcac	aaggataccg	cttcttacia	ccttaccctg	gcttatgctc	cagata
2401	aagaagcaac	cctgactgta	ctgcctctct	gttagtgagc	ccaacctctg	ctgtct
2461	aactaaagct	aataaccttg	cgcggcatgc	tttcatatta	caagcaggaa	attact
2521	tttaactcgc	aacacggagc	tcttcagtca	attcggtttc	gaactccgag	gttctt
2581	cacctataac	atagatctcg	gatcgaagat	ccagttctaa	tctcatccac	ctcccc
2641	ccgtagacac	ggagcagggc	ctccctttgt	ctactctagc	tggtgacctt	gacttt
2701	tatttttatg	atcgagttag	gatatctcaa	atcacccaaa	gcattgaattt	atgaaa
2761	cagtctactg	gttcttaata	tcttcgagcc	tatttgctc	gaattctttg	agcttc
2821	acgacgctca	aacagcctta	actccctccg	atagctataa	tggaaatgtg	acctct
2881	agttccaggt	aaaagaaact	tcatacaggaa	caacgtatac	ttgtgaaggc	aatgtg
2941	tctcctttgc	agggaaagat	tcaggtctaa	agaaaagttg	tttctcagct	actgat
3001	ttaccttcct	aggaaacggg	tatactcttt	gctttgataa	tattactact	acagct
3061	accccgagc	cattaatgtt	caaggtcaag	gaaaaacctt	aggcatctca	ggattt
3121	tattttcatg	tgcttattgt	cctccaggca	caactgggtta	cggagctata	cagact
3181	gcaacacaac	tttaaaagat	aactctagtc	ttgtcttcca	taaaaactgc	tcaaca
3241	aaggtggggc	tatccagtgt	aaaggaagca	gtgatgctga	attaaaaata	gaaaat
3301	agaatctgg	tttctcagaa	aactcctcca	cttcaaaagg	cggggctatt	tatgct
3361	aactcaccat	tgtctcaggt	gggcctacat	tattttctaa	caactctgta	tccaac
3421	catcccctaa	aggcggagct	attagcataa	aagattcaag	tggtgaatgt	agccta
3481	ctgatctcgg	agatattacc	ttcgatggga	acaaaatcat	caaaactagt	ggtgga
3541	ctacagtaac	aagaaattcc	atagatctcg	gcacagggaa	atttacaag	ctacgt
3601	aagacggctt	cgaatttttc	ttctatgacc	ctattactgg	gggaggatct	gatgaa
3661	acattaataa	aaaagaaact	gttgattata	caggaaagat	cgtcttctca	ggtgaa
3721	tatccgatga	agaaaaagca	cgagcggaaa	acctagcttc	tactttcaac	caaccc
3781	cattatcagc	aggatctctt	gtacttaaag	atgggtgtatc	tgtaacgcga	aaacaa
3841	cgcaggaagc	gggatctacc	gttgatcatg	atctagggac	cacattacag	acgcct
3901	caggtggaga	aaccatcacc	ctaactaatc	tagatattaa	catcgctcgc	ttgggg
3961	gggggggtac	ctctcctgct	aaactcgcaa	caaatacagc	aagtcaagct	ataact
4021	acgctgtcaa	tctagtcgat	gctgatggca	atgcttatga	agatcctatt	cttgct
4081	ctaaaccttt	cacagcaata	gtagctacaa	ctaacgctag	tacagtcaca	cagcct
4141	ataatctaac	aaattatgtc	cctcctactc	attacgggtta	ccaaggaaat	tggaca
4201	cttgggacac	cgaacacagc	acaaaaacag	ccactctaac	ttgggaacaa	actggc
4261	cccctaacc	agaacgtcaa	ggacctttag	tcccgaatac	tctttgggg	gcattc
4321	acctcagagc	tatacaaaac	ttaatggata	ttagcgtcaa	tggcgtgac	taccat
4381	gtttttgggt	atccgggtcta	gctaacttct	tacacaaaag	tggctctgat	actaaa
4441	agttccgtca	caatagcgcc	ggatacgctt	taggcgtcta	cgcacaaaact	ccttct
4501	atattttcag	tgcggctttc	tgccaactct	tcggaaagga	caaagactat	ttagtg

4561 aaaacaacgc caacattttac gcaggtttctc tctattatca gcatatctcc tattgg  
4621 cttggcagaa tctgctacaa aacactatcg gtgcagaagc tccgttagtc cttaac  
4681 agttaactta ttgtcatgct tcaaacgaca tgaaaaccaa catgacgact acttac  
4741 ctcgtaaaac aacgtatgca gaaatcaagg gtgattgggg taacgattgt ttcgga  
4801 agcttggtgc aactgtgcct atccaaacag aatctttctc cctattcgat atgtac  
4861 ctttcctgaa gtttcaactt gtgcatacgc accaagatga ctttaaggaa aacaat  
4921 atcaggggaag atacttcgaa agcagcaatc tcaccaacct ttctctgcct atcggc  
4981 agtttgagag atttgctaac aacgatacag cttcttatca tgtcactgct gcttat  
5041 ctgatatcgt aagaagtaac cctgactgta ctacttctct gttagtaagc cccgac  
5101 ctgtctgggt aacgaaagcc aacaaccttg cgcgaagcgc cttcatgcta caagca  
5161 actacttgtc tttaagtcac aacatagaaa tcttcagcca gttcggtttc gagctc  
5221 gatcttcacg aacctataac gtagatctcg gatcgaagat ccagttctaa tctcat  
5281 ctctcctacc ccgttcccac ggggtagggc cttataacct tgactctcct tggaat  
5341 cgaacaaaga gtcaaaaaaa tacgtctcat aggattctgg tctattccct tttctt  
5401 ttggagaaaa atctgatcac ctgcaagaca cgcaagatat aggtaaattt cgtcaa  
5461 ccgaatcggt aaaaccttga tcttctagaa gttttttatt ttagatggtc atctaa  
5521 aagataggcc tcggcatttt cctcatcaag atacagcaac tcttgcttta ttttcc  
5581 tcccttatgt caggtagagc cgcaaaatat gcagggaata ctaatgaaaa actcta  
5641 tggggtttta ctgttttcct cttttgcctt atccactgct accaaacttc ttgcag  
5701 cgactctgtc aaccttgcaa ctggattcaa cggctccact agtgaaactt tcaatg  
5761 acaaacagat aatgctgacg ggacaacata tattctaggc agcgcgatca cctttg  
5821 cataaatcaa ttaaaaccag caaacactag ctgttttgct aatacagctg gagatc  
5881 gtttactggg aatcgacgac ttctctatatt caataatatt tcatcaacag cgaaag  
5941 cgctatcagc acaactgcgg atggtaagac actcacaata tccggggctc tacaac  
6001 tttctacatg tcgccaagat tggccacggg aaatggcgctc atttattcta atagct  
6061 actcatcgag aacaattctc aaggtagctc gggactgaat aagtctgcag

//

[Disclaimer](#) | [Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)

Mar 17 2003 10:55:57



PubMed	Nucleotide	Protein	Genome	Structure	PMC	Taxonomy	OMI
Search <u>Protein</u> <input type="button" value="v"/> for <input type="text"/>							
		Limits	Preview/Index	History	Clipboard	D	
Display	default	<input type="button" value="v"/> Show: 20	<input type="button" value="v"/> Send to	File	<input type="button" value="v"/>	Get Subsequence	

☐ 1: AAC15922.POMP90A precursor...[gi:1617507]

BLink, Domains, Links

LOCUS AAC15922 839 aa linear BCT 07-

DEFINITION POMP90A precursor [Chlamydophila abortus].

ACCESSION AAC15922

VERSION AAC15922.1 GI:1617507

DBSOURCE locus CPU65942 accession U65942.1

KEYWORDS .

SOURCE Chlamydophila abortus

ORGANISM Chlamydophila abortus

Bacteria; Chlamydiae; Chlamydiales; Chlamydiaceae; Chlamydo

REFERENCE 1 (sites)

AUTHORS Longbottom,D., Russell,M., Jones,G.E., Lainson,F.A. and Herring,A.J.

TITLE Identification of a multigene family coding for the 90 kDa of the ovine abortion subtype of Chlamydia psittaci

JOURNAL FEMS Microbiol. Lett. 142 (2-3), 277-281 (1996)

MEDLINE 96406378

PUBMED 8810511

REFERENCE 2 (sites)

AUTHORS Longbottom,D., Russell,M., Dunbar,S.M., Jones,G.E. and Herr

TITLE Molecular cloning and characterization of the genes coding highly immunogenic cluster of 90-kilodalton envelope protei the Chlamydia psittaci subtype that causes abortion in shee

JOURNAL Infect. Immun. 66 (4), 1317-1324 (1998)

MEDLINE 98187897

PUBMED 9529048

REFERENCE 3 (residues 1 to 839)

AUTHORS Longbottom,D.

TITLE Direct Submission

JOURNAL Submitted (02-AUG-1996) Moredun Research Institute, Interna Research Centre, Pentlands Science Park, Bush Loan, Penicui Midlothian EH26 0PZ, UK

FEATURES Location/Qualifiers

source 1..839

/organism="Chlamydophila abortus"

/strain="S26/3"

/specific\_host="Ovis aries"

/db\_xref="taxon:83555"

/clone="S112"

/clone\_lib="lambda EMBL-3"

Protein /dev\_stage="elementary body"  
/note="submitted as Chlamydia psittaci"  
1..839  
/product="POMP90A precursor"  
sig peptide 1..16  
/note="putative"  
mat peptide 17..838  
/product="POMP90A"  
/note="putative"  
CDS 1..839  
/coded\_by="U65942.1:2751..5270"  
/note="encodes putative 89,836 Da outer membrane p  
/citation=[2]  
/transl\_table=11


## ORIGIN

```
1 mkhpyvwfli ssslfasnsl sfandaqtal tpsdsyngnv tseefqvket ssgtty
61 nvcisfagkd sglkkscfsa tdnltflgng ytlcfdnitt tasnpgainv qgggkt
121 gfslfscayc ppgttgygai qtkgnttlkd nsslvfhknc staeggaiqc kgssda
181 ennqnlvfse nsstskggai yadklktivsg gptlfsnnsv sngsspkgga isikds
241 sltadlgdit fdgnkiipts ggsstvttrns idlgtgkftk lrakdgfgif fydpti
301 delninkket vdytgkivfs geklsdeeka raenlastfn qpitlsagsl vlkdgv
361 kqvtqeagst vvmldgttlq tpssggetit ltnldinias lggggggtspa klatnt
421 itinavnlvd adgnayedpi latskpftai vattnastvt qptdnltnyv ppthyg
481 wtvtdwteta tktatltweq tgyspnperq gplvpntlwg afsdlraiqn lmdisv
541 yhrqfwvsgl anflhksgsd tkrkfrhnsa gyalgvyakt psddifsaaf cqlfgk
601 lvsknnaniy agslyyqhis ywsawqnllq ntigaeaplv lnaqltycha sndmkt
661 tyaprkttya eikgdwgnlc fgvelgatvp iqtessllfd myspflkfql vhthqd
721 nnsdqgryfe ssnltnlslp igikferfan ndtasyhvta ayspdivrsn pdctts
781 pdsavwvтка nnlarsafml qagnylslsh nieifsqfgf elrgssrtyv vdlgsk
```

//

Disclaimer | Write to the Help Desk  
NCBI | NLM | NIH

Mar 17 2003 10:55:57



PubMed	Nucleotide	Protein	Genome	Structure	PMC	Taxonomy	OMI
Search <input type="text" value="Protein"/> for <input type="text"/>							
Limits		Preview/Index		History		Clipboard	
Display	<input type="text" value="default"/>	Show:	<input type="text" value="20"/>	Send to	<input type="text" value="File"/>	Get Subsequence	

☐ 1: AAC15924. POMP90B precursor...[gi:1617510]

BLink, Domains, Links

LOCUS AAC15924 839 aa linear BCT 07-

DEFINITION POMP90B precursor [Chlamydophila abortus].

ACCESSION AAC15924

VERSION AAC15924.1 GI:1617510

DBSOURCE locus CPU65943 accession U65943.1

KEYWORDS .

SOURCE Chlamydophila abortus

ORGANISM Chlamydophila abortus

Bacteria; Chlamydiae; Chlamydiales; Chlamydiaceae; Chlamydo

REFERENCE 1 (sites)

AUTHORS Longbottom,D., Russell,M., Jones,G.E., Lainson,F.A. and Herring,A.J.

TITLE Identification of a multigene family coding for the 90 kDa of the ovine abortion subtype of Chlamydia psittaci

JOURNAL FEMS Microbiol. Lett. 142 (2-3), 277-281 (1996)

MEDLINE 96406378

PUBMED 8810511

REFERENCE 2 (sites)

AUTHORS Longbottom,D., Russell,M., Dunbar,S.M., Jones,G.E. and Herr

TITLE Molecular cloning and characterization of the genes coding highly immunogenic cluster of 90-kilodalton envelope protei the Chlamydia psittaci subtype that causes abortion in shee

JOURNAL Infect. Immun. 66 (4), 1317-1324 (1998)

MEDLINE 98187897

PUBMED 9529048

REFERENCE 3 (residues 1 to 839)

AUTHORS Longbottom,D.

TITLE Direct Submission

JOURNAL Submitted (02-AUG-1996) Moredun Research Institute, Interna Research Centre, Pentlands Science Park, Bush Loan, Penicui Midlothian EH26 0PZ, UK

FEATURES Location/Qualifiers

source 1..839

/organism="Chlamydophila abortus"

/strain="S26/3"

/specific\_host="Ovis aries"

/db\_xref="taxon:83555"

/clone="S21 and S432"

/clone\_lib="lambda EMBL-3"

Protein /dev\_stage="elementary body"  
/note="submitted as Chlamydia psittaci"  
1..839  
/product="POMP90B precursor"  
sig peptide 1..16  
/note="putative"  
mat peptide 17..839  
/product="POMP90B"  
/note="putative"  
CDS 1..839  
/coded\_by="U65943.1:3290..5809"  
/note="encodes putative 89,836 Da outer membrane p  
/citation=[2]  
/transl\_table=11

## ORIGIN

```
1 mkhpyvwfli ssslfasnsf sfandaqtal tpsdsyngnv tseefqvket ssgtty
61 nvcisfagkd sglkkscfsa tdnltflgng ytlcfdnitt tasnpgainv qgggkt
121 gfslfscayc ppgttgylgai qtkgnttlkd nsslvfhknc staeggaiqc kgssda
181 ennqnlvfse nsstskggai yadkltivsg gptlfsnnsv sngsspkgga isikds
241 sltadlgdit fdgnkiikts ggsstvttrns idlgtgkftk lrakdgfgif fydpit
301 delninkket vdytgkivfs geklsdeeka raenlastfn qpitlsagsl vlkdgv
361 kqvtqeagst vvmldgttlq tpssggetit ltnldinias lggggggtspa klatnt
421 itinavnlvd adgnayedpi latskpftai vattnastvt qptdnltnyv ppthyg
481 wtvtdwteta tktatltweq tgyspnperq gplvpntlwg afsdlraiqn lmdisv
541 yhrqfwvsgl anflhksgsd tkrkfrhnsa gyalgvyakt psddifsaaf cqlfgk
601 lvsknnaniy agslyyqhis ywsawqnlq ntigaeaplv lnaqltycha sndmkt
661 tyaprkttya eikgdwgnhc fgvelgatvp iqtessllfd myspflkfql vhthqd
721 nnsdqgryfe ssnltnslslp igikferfan ndtasyhvta ayspdivrsn pdctts
781 pdsavwvтка nnlarsafml qagnylslsh nieifsqfgf elrgssrtyv vdlgsk
```

//

Disclaimer | Write to the Help Desk  
NCBI | NLM | NIH

Mar 17 2003 10:55:57

RESULT 2

P77792

ID P77792 PRELIMINARY; PRT; 839 AA.  
 AC P77792;  
 DT 01-FEB-1997 (TrEMBLrel. 02, Created)  
 DT 01-FEB-1997 (TrEMBLrel. 02, Last sequence update)  
 DT 01-JUN-2002 (TrEMBLrel. 21, Last annotation update)  
 DE POMP90B precursor.  
 OS Chlamydia psittaci (Chlamydophila psittaci).  
 OC Bacteria; Chlamydiales; Chlamydiaceae; Chlamydophila.  
 OX NCBI\_TaxID=83554;  
 RN [1]  
 RP SEQUENCE FROM N.A.  
 RC STRAIN=OVINE ENZOOTIC ABORTION ISOLATE S26/3;  
 RA Longbottom D., Russell M., Jones G.E., Lainson F.A., Herring A.J.;  
 RL Submitted (OCT-1996) to the EMBL/GenBank/DDBJ databases.  
 RN [2]  
 RP SEQUENCE FROM N.A.  
 RC STRAIN=OVINE ENZOOTIC ABORTION ISOLATE S26/3;  
 RA Longbottom D., Russell M., Dunbar S.M., Jones G.E., Herring A.J.;  
 RL Submitted (OCT-1996) to the EMBL/GenBank/DDBJ databases.  
 DR EMBL; U65943; AAC15924.1; -.  
 DR EMBL; U65942; AAC15922.1; -.  
 DR InterPro; IPR003368; Chlamydia\_PMP.  
 DR InterPro; IPR003357; OMP.  
 DR Pfam; PF02415; DUF145; 1.  
 DR Pfam; PF02385; OMP; 1.  
 KW Signal.  
 FT SIGNAL 1 16 POTENTIAL.  
 FT CHAIN 17 839 POMP90B.  
 SQ SEQUENCE 839 AA; 89824 MW; 4581C7CBAF7FF4C4 CRC64;

Query Match 1.3%; Score 12; DB 2; Length 839;  
 Best Local Similarity 100.0%; Pred. No. 0.013;  
 Matches 12; Conservative 0; Mismatches 0; Indels 0; Gaps 0;

QY 602 PNPERQGPLVPN 613  
 |||||  
 Db 505 PNPERQGPLVPN 516

## Molecular Cloning and Characterization of the Genes Coding for the Highly Immunogenic Cluster of 90-Kilodalton Envelope Proteins from the *Chlamydia psittaci* Subtype That Causes Abortion in Sheep

DAVID LONGBOTTOM,\* MARY RUSSELL,† SUSANNA M. DUNBAR, GARETH E. JONES,  
AND ALAN J. HERRING‡

Moredun Research Institute, Edinburgh EH17 7JH, United Kingdom

Received 11 July 1997/Returned for modification 10 September 1997/Accepted 9 January 1998

Proteins present in the outer membrane of chlamydiae that are involved in mucosal epithelial cell infection must clearly be identified and characterized if we are to understand and modify the pathogenic mechanisms utilized by these organisms. We have identified and isolated a family of four genes encoding putative outer membrane proteins (POMPs), a group of proteins of approximately 90 kDa present in the outer membrane of the subtype of *Chlamydia psittaci* that causes ovine enzootic abortion (strain S26/3). These proteins, although minor components, are major immunogens, as shown by the immunoblotting of chlamydial outer membrane complexes with postabortion sheep sera, and are therefore potential diagnostic and/or protective antigen candidates. Immunoblotting of the expressed amino- and carboxy-terminal halves of one of the POMPs with postabortion sheep sera showed that the major humoral immune response appeared to be directed solely against the amino-terminal half. This result, in combination with the positive immunofluorescence staining of S26/3-infected cells using POMP-specific (specific to the amino-terminal half of the proteins) monoclonal antibodies, suggests the probable surface localization of the POMPs and, more specifically, the surface exposure of the amino-terminal half of these proteins. The four *pomp* genes are highly homologous, sharing 82 to 100% similarity with each other (two of the genes are identical). Genes with strong and weak homologies were also detected in *C. psittaci* avian and feline pneumonitis strains, respectively. No *pomp* homologs were found in strains of *C. trachomatis* and *C. pneumoniae*, but this does not preclude their existence. The absence of homology with various subtypes of *C. pecorum*, which complicate the diagnosis of the ovine abortion subtype, indicates the possible suitability of these 90-kDa proteins as serodiagnostic antigens.

The elementary body (EB) outer cell envelope of members of the genus *Chlamydia* resembles that of gram-negative bacteria in that it contains lipopolysaccharide but is atypical in lacking a peptidoglycan layer (9). The structural stability of EBs is attributed to extensive intermolecular disulfide bond cross-linking of the 40-kDa major outer membrane protein (MOMP), one or two large (60-kDa), cysteine-rich proteins (EnvB or Omp2), and a small (12-kDa), cysteine-rich lipoprotein (EnvA or Omp3) forming a supramacromolecular lattice (reviewed in reference 25). While these are the major protein components of the outer membrane, other minor and so far uncharacterized proteins have been identified which may have major functional roles (25, 26).

Since the EB outer surface has such an important role in the infection process, it is not surprising that its outer membrane has been the focus of much research, particularly with regard to serodiagnosis and vaccine design. Currently, our research is focused on trying to improve serodiagnosis (2) and developing a recombinant vaccine (15) against the subtype of *Chlamydia*

*psittaci* which causes ovine enzootic abortion (OEA). This infection is the most common infectious cause of abortion in sheep in the United Kingdom (UK) (1) and can also cause abortion in women (13). We have been particularly interested in protein antigens located in the outer membrane of OEA *C. psittaci* which may be involved in protective immunity. The only protein consistently shown to be exposed on the surface of the EB is the MOMP (3, 4, 6, 29). It is the major component of the chlamydial outer membrane complex (COMC), which we have previously shown protects pregnant ewes against experimentally induced OEA (30).

In addition to the MOMP, the *C. psittaci* COMC contains a group of minor proteins with sizes of approximately 90 kDa which have been shown to react very strongly in immunoblotting experiments (5, 12, 19, 27) and to be important serodiagnostic antigen candidates (27). Recently, we have reported the isolation of two recombinant  $\lambda$ gt11 clones (named 31 and 99) which contain fragments of the genes coding for these 90-kDa proteins, as well as the finding that these proteins belong to a multigene family (18). This report describes the isolation and characterization of four genes which belong to this family and provides evidence of the location and possible orientation of their protein products in the outer membrane of *C. psittaci*.

### MATERIALS AND METHODS

**Chlamydia.** OEA *C. psittaci* S26/3 was grown in L929 cells, and *C. pecorum* strains were cultured in embryonated chicken eggs (20). EBs were purified from infected cells or chicken embryo yolk sacs by density gradient centrifugation, and genomic DNA was extracted as described previously (20). In addition to the *C. psittaci* OEA strain, genomic DNAs from the following *Chlamydia* strains were

\* Corresponding author. Mailing address: Moredun Research Institute, International Research Centre, Pentlands Science Park, Bush Loan, Penicuik, Midlothian EH26 OP2, United Kingdom. Phone: 44 (0)131 664 3262. Fax: 44 (0)131 664 8001. E-mail: longd@mri.sari.ac.uk.

† Present address: Department of Structural Biochemistry, University of Edinburgh, Edinburgh EH9 3JR, United Kingdom.

‡ Present address: The Public Health Laboratory, Kingsdown, Bristol BS2 8EL, United Kingdom.



also studied: (i) *C. psittaci* avian (cockatiel isolate 725; described in reference 20) and feline pneumonitis (FePn; described in reference 21) isolates; (ii) *C. pecorum* subtypes P787 (causing ovine arthritis), W73 (causing ovine enteritis), PV3056/3 (causing bovine metritis), and JP1751 (ovine normal feces); (iii) *C. trachomatis* (VW1 serovar D; DNA obtained from J. Trehanne, Institute of Ophthalmology, London, UK); and (iv) *C. pneumoniae* (IOL-207; DNA obtained from J. Trehanne).

For immunofluorescence microscopy, McCoy cells were grown on sterile glass coverslips in tissue bottles (Bibby Sterilin Ltd., Stone, UK) and infected with OEA *C. psittaci* (inoculum titer of  $5 \times 10^8$  inclusion-forming units/ml) as previously described (20).

**Hybridization screening.** A partially *MhoI*-digested S26/3 genomic DNA library constructed in lambda replacement vector EMBL-3 (Stratagene Ltd., Cambridge, UK) from S26/3 DNA (14) was screened by standard techniques using digoxigenin (DIG)-labelled *p90f31* (216-bp fragment) and *p91Bf99* (450-bp *EcoRI* fragment) probes (clone 31 and 99 DNA inserts, respectively; described in reference 18). Recombinant phage were plated by using the restrictive host strain *Escherichia coli* Q359. Recombinant lambda DNA was isolated from hybridization-positive clones by plate lysates using the permissive strain *E. coli* LE392 and purified from lambda phage particles by polyethylene glycol 6000 precipitation in accordance with standard procedures.

**Sequencing and gene analysis.** The recombinant lambda DNA clones were Southern blotted by using *p90f31* and *p91Bf99* DIG-labelled probes, and probe-specific restriction enzyme fragments were cloned into pBluescript (Stratagene) as described previously (18). PCR amplification of recombinant lambda DNA was performed by using the Expand Long Template PCR System (Boehringer Mannheim, Lewes, UK) and primers which flanked the cloning site of EMBL-3 (5'-GCGCACTCGTGAAGGAGTAGGCGGA-3' and 5'-TGAACTCGTCGCGAATAACGAGTGGAT-3') and those based on sequence information obtained from the cloned restriction enzyme fragments. PCR products were subcloned into vector pGEM-T (Promega, Southampton, UK) as previously described (18). Recombinant plasmid DNA was purified by using a QIAprep Spin Plasmid Miniprep Kit (QIAGEN Ltd., Crawley, UK) for manual sequencing by the dideoxy-chain termination method using the Sequenase Version 2.0 DNA Sequencing Kit (Amersham International plc, Little Chalfont, UK) and using a QIAGEN Plasmid Kit (QIAGEN Ltd.) for automated Taq FS cycle sequencing (at The Advanced Biotechnology Centre, The Charing Cross and Westminster Medical School, London, UK). All internal primers were synthesized by Oswel DNA Service (Medical and Biological Sciences Building, University of Southampton, Southampton, UK). Recombinant plasmid DNA was sequenced manually at least once in each direction. All sequencing problems, primarily due to secondary structure, were resolved by automated cycle sequencing.

DNA and deduced amino acid sequences were analyzed by using the Wisconsin GCG software package on Seqnet (Daresbury Laboratory) and DNASTAR software (DNASTAR Inc.). The protein sequences were also analyzed by using the program PSORT (on the World Wide Web at URL <http://psort.nibb.ac.jp>) (23), which predicts the presence of signal sequences by the methods of McGeech (22) and von Heijne (31) and detects potential transmembrane domains by the method of Klein et al. (17). DNA and amino acid sequences were compared with those in the GenBank/EMBL and SwissProt databases by using the GCG FASTA program and the Genome Sequence Database (GSDB) (on the World Wide Web at URL <http://www.tigr.org>) (8, 10).

**PCR analysis.** *C. psittaci* S26/3 genomic DNA fragments containing the four *pomp* genes were amplified by PCR by using the Expand Long Template PCR System and mutually exclusive *pomp* flanking region primers: primer pairs S5766 (5'-GTAAGCAAGAACGCCGCAACG-3') plus P0787 (5'-CTAGTGGAGC CGTTGAATTCAGTTGC-3') and S5765 (5'-TGAGCCAAATCCTTCTACAG CAAT-3') plus S2429 (5'-AGTCTTAGCGTCACAGGCCGAG-3') for regions A and B, respectively (see Fig. 2). PCR products were analyzed by restriction endonuclease digestion (*EcoRI*, *PstI*, and *XbaI*; single and double digests).

**Immunoblotting.** *C. psittaci* S26/3 EBs were purified from infected L929 cells by density gradient centrifugation as described previously (20). COMCs prepared from EBs as the Sarkosyl-dithiothreitol (DTT)-insoluble fraction (30) were solubilized in *n*-octyl- $\alpha$ -D-glucopyranoside-DTT (19) and subjected to sodium dodecyl sulfate (SDS)-polyacrylamide gel electrophoresis (PAGE) on 7.5% gels after boiling in sample buffer containing 2% SDS and 5% 2-mercaptoethanol. Protein was transferred to a nitrocellulose membrane and incubated with P90F31 and P91Bf99 (protein products of gene fragments *p90f31* and *p91Bf99*, respectively) affinity-purified antibodies or a pool of postabortion sera from 10 ewes experimentally infected with OEA *C. psittaci*, and antibody binding was detected as described previously (18).

**Expression analysis.** The PCR-generated 5' and 3' halves of *pomp90A* were cloned in frame into expression vector pET-22b(+) (Novagen, R&D Systems Europe Ltd., Abingdon, UK) via engineered *NdeI*/*HindIII* and *NdeI*/*EcoRI* restriction enzyme sites, respectively, in the primers by using procedures described previously (18). The 1.16-kb 5' and 1.26-kb 3' fragments were generated by using primer pairs P0786 (5'-CCAAAGCATGAACATATGAAACATCCAGTC-3') plus V8704 (5'-AAGAAAGCTTCTGTATGTGGTCCCTAG-3') and T3252 (5'-AACGCTGTCCATATGGTCTGATGCGGC-3') plus V8703 (5'-GAGGTGGATGAGAATCAACTGGATCTTCG-3'), respectively. Initial cloning was performed in *recA* mutant strain JM109, which lacks the gene for T7 RNA

polymerase, to allow examination of the construct sequences. The pET constructs were subsequently transformed into expression host BL21(DE3), which contains the T7 RNA polymerase gene, and induced at 37°C for 3 h with isopropyl- $\beta$ -D-thiogalactopyranoside (IPTG) at a final concentration of 1 mM. Expressed products were purified from inclusion bodies by a differential centrifugation procedure as detailed in Novagen's pET System Manual (24) and analyzed by Western blotting as described in the paragraph on immunoblotting but also by using 12 anti-90-kDa protein monoclonal antibodies raised against whole EBs from *C. psittaci* abortion strains (three monoclonal antibodies) and a parakeet strain (nine monoclonal antibodies). All of these monoclonal antibodies were obtained from E. Vretou, Institut Pasteur Hellenique, Athens, Greece (the original source of the avian monoclonal antibodies was W. Schuy, Behringwerke, Marburg, Germany), and have been described previously (32). In particular, the avian strain monoclonal antibodies are known to cross-react with the ovine abortion strain 90-kDa antigen triplet. A summary of the various antibodies used in this study is shown in Table 1.

**Immunizations.** Two 500- $\mu$ g doses of the purified carboxyl half of POMP90A in adjuvant Montanide ISA 773 (SEPPIC, Paris, France) were administered subcutaneously 2 weeks apart to four hogs (1- to 2-year-old sheep). All animals were bled pre-vaccination and 3 weeks after the second vaccination. Sera were analyzed by Western blotting of whole EB preparations as described previously (18).

**Immunofluorescence.** OEA strain-infected McCoy cells, grown on coverslips for various times (18 to 72 h) postinfection, were briefly washed in phosphate-buffered saline (PBS) and fixed with acetone for 10 min. After air drying for 15 min, coverslips were stored at -70°C. Upon removal from the freezer, coverslips were washed in PBS and incubated with 2% bovine serum albumin in PBS for 30 min and then incubated for 1 h at 37°C in primary antibody diluted in 2% bovine serum albumin-PBS. Coverslips were washed three times for 5 min each time with PBS and then incubated for 1 h at 37°C in fluorescein-conjugated sheep anti-mouse secondary antibody (Scottish Antibody Production Unit, Law Hospital, Carlisle, Scotland, United Kingdom) diluted 1/80 in PBS. After a further three washes, coverslips were inverted onto a drop of Citifluor (glycerol-PBS mixture) (Citifluor, Canterbury, UK) and examined under an Olympus BX50 microscope using a 40 $\times$  objective.

**Southern blot analysis.** Genomic DNA (500-ng aliquots) prepared from *C. psittaci* cockatiel and FePn isolates, as well as *C. trachomatis* (VW1 serovar D), *C. pneumoniae* (IOL-207), and various *C. pecorum* (P787, W73, PV3056/3, and JP1751) subtypes were Southern blotted by using DIG-labelled *p91Bf99* (450-bp *EcoRI* fragment) and *pomp90A* (1.1-kb *EcoRI*/*XbaI* and 1.6-kb *XbaI* fragments) probes (probe concentration of 25 ng/ml) as previously described (18). The hybridization buffer consisted of 25% deionized formamide, 5 $\times$  SSC (1 $\times$  SSC is 0.15 M NaCl plus 0.015 M sodium citrate), 0.02% SDS, and 2% blocking reagent (Boehringer Mannheim). Hybridization was performed at 37°C for 17 h. The most stringent wash was in 2 $\times$  SSC-0.1% SDS at 37°C. Detection of probe binding was with the chemiluminescent substrate CSPD (Boehringer Mannheim).

**Nucleotide sequence accession numbers.** The DNA sequences reported here were submitted to the GenBank/EMBL databases and assigned accession no. U65942 and U65943.

## RESULTS

**Cloning and sequence analysis of the genes for the 90-kDa proteins.** Five positive clones were isolated from the  $\lambda$ EMBL-3 library, and restriction endonuclease-digested recombinant lambda DNA was analyzed by Southern blotting by using *p90f31* (data not shown) and *p91Bf99* (Fig. 1) probes. All of the fragments previously observed by Southern blotting of digests of S26/3 genomic DNA with these probes (18) were identified. Initially, the 0.45-, 3.7-, 4.0-, and 5.3-kb *EcoRI* fragments were subcloned and sequenced. Further sequence analysis was performed on PCR-derived DNA fragments, although ultimately all of the sequence information presented here was obtained from  $\lambda$ EMBL-3 clone-derived restriction enzyme fragments. The combination of information obtained from the Southern blotting, mapping, and sequencing of these EMBL-3 clones enabled the maps shown in Fig. 2 to be constructed. The maps were verified further by the restriction enzyme analysis of PCR-generated 6.5- and 5.6-kb DNA fragments of the two OEA strain genomic DNA regions (A and B in Fig. 2) containing the *pomp91A* and *pomp90A* and the *pomp91B* and *pomp90B* genes, respectively (results not shown).

Sequence analysis identified four possible open reading frames (ORFs) designated as follows: *pomp* for putative outer membrane protein, 90 or 91 for the predicted molecular mass

TABLE 1. Antibody reagents used in this study<sup>a</sup>

Antibody	Antibody type	Antigen source	Antigen recognition	Source or reference
181 191 192	Ascites	<i>C. psittaci</i> abortion strain EBs	90-kDa triplet; 49-kDa amino-terminal fragment of POMP90A	32
89-79/0040  89-84/019  89-76/073 89-76/049 89-76/030  89-73/053  89-73/04 89-73/0200 89-73/0248	Tissue culture supernatant	<i>C. psittaci</i> avian (parakeet) strain EBs	90-kDa triplet; P91BF99  90-kDa triplet; P90F31  90-kDa triplet; 49-kDa amino-terminal fragment of POMP90A	W. Schuy, Behringwerke, Marburg, Germany; 32
Anti-P90F31	Affinity-purified monospecific antiserum	P90F31	90-kDa triplet; P90F31	18
Anti-P91BF99		P91BF99	90-kDa triplet; P91BF99	

<sup>a</sup> Antibodies 181, 191, and 192 are serotype 1 specific (32). Either mouse monoclonal antibodies or antibodies affinity purified by using pooled postabortion sera from ewes experimentally infected with OEA *C. psittaci* were used. The antigen source for antibody production or purification was either formalin-inactivated whole EBs or SDS-PAGE-denatured antigen.

(in kilodaltons) of the translated product, and *A* or *B* to distinguish the two different locations of these genes in the chlamydial genome (Fig. 2). Genes *pomp90A* and *pomp90B* are identical, each having an ORF of 2,520 bp encoding a protein (POMP90A and POMP90B, respectively) with a predicted molecular mass of 89,836 Da and pI 5.27. Gene *pomp91A* has an ORF of 2,544 bp encoding a protein (POMP91A) with a predicted mass of 90,707 Da and pI 5.45, and *pomp91B* has an

ORF of 2,541 bp encoding a protein (POMP91B) with a predicted mass of 90,847 Da and pI 6.08. These predicted molecular masses and pI values correlate well with those determined by two-dimensional electrophoretic analysis (33). Overall, the *pomp91A* and *pomp91B* sequences are 85% similar and both are 82% similar to that of *pomp90A* or *pomp90B*. Consensus promoter sequences, ribosomal binding sites, and transcription termination sequences could be identified for each gene (see GenBank database entries U65942 and U65943). In addition, polyguanine tracts, consisting of 13 to 15 dGTP groups, could be found located centrally in each of the four genes (Fig. 3). Analysis of the sequences also confirmed our previous suggestion that *p90f31* and *p91bf99* were derived from different genes (18), with *p90f31* originating from either *pomp90A* or *pomp90B* and *p91bf99* originating from *pomp91B* (Fig. 2). Searches in the GenBank/EMBL, GSDB, SwissProt databases failed to find any related DNA or protein sequences.

Figure 4 shows a multiple sequence alignment of the four proteins and highlights the similarity of the sequences. POMP91A and POMP91B were found to be 89% similar to each other and 86% and 87% similar, respectively, to POMP90A and POMP90B. Analysis of the N-terminal protein sequences by the PSORT protein localization prediction program showed that all of the POMP's have predicted signal sequences and, with the exception of POMP91A, signal peptidase I cleavage sites (Fig. 4). Furthermore, no hydrophobic transmembrane regions were identified and the overall prediction, again with the exception of POMP91A, was that the proteins are located in the outer membrane. This correlated with secondary-structure analyses (11) predicting that they consist almost entirely of  $\beta$ -sheet, a normal conformation for bacterial outer membrane proteins where the membrane-spanning regions are formed by the  $\beta$  strands. In addition, the last




FIG. 1. Southern blot analysis of ovine abortion strain S26/3  $\lambda$ EMBL-3 clone DNA. Aliquots (50 ng) of recombinant lambda DNA prepared from clones S432 (lanes 1 and 2), S112 (lanes 3 to 5), and S21 (lanes 6 to 8) were digested with *EcoRI* (lanes 1, 3, and 6), *PstI* (lanes 2, 4, and 7), and *XbaI* (lanes 5 and 8) and Southern blotted by using a DIG-labelled *p91bf99* probe as described in Materials and Methods. DNA size markers are indicated on the left.




FIG. 2. Restriction enzyme maps for the family of 90-kDa *pomp* genes. The positions of the restriction enzyme sites (expressed in base pairs), deduced from the sequencing (hatched areas) and Southern blotting results, are in parentheses. A and B represent two separate regions of the OEA *C. psittaci* genome. The sequences encoded by  $\lambda$ EMBL-3 clones S81, S112, and S421 for region A (A) and by clones S21 and S432 for region B (B) are as illustrated. The sequences incorporating *pomp91A* and *pomp90A* (positions 1 to 6110 in A) and *pomp91B* and *pomp90B* (positions 1 to 6234 in B) have been submitted to the GenBank/EMBL databases and assigned accession no. U65942 and U65943, respectively. The double-headed arrow depicts the region of complete identity in A (positions 2111 to 5312) and B (positions 2650 to 5851). The *pomp* flanking region forward primers S5766 (P1) and S5765 (P3) and reverse primers P0787 (P2) and S2429 (P4) bind at the indicated positions.

residue of all four proteins is phenylalanine, a feature found in the majority of bacterial outer membrane proteins (28).

**Immunoblotting analysis of COMCs.** When COMCs were immunoblotted by using a pool of postabortion sera from experimentally infected ewes, the major reaction was with the MOMP, which comprises about 90% of the protein detected by Coomassie blue staining and typically occurs in these preparations as a double band of about 38 and 36 kDa. However, a triplet of bands with molecular masses in the region of 90 to 95 kDa was also clearly recognized, together with proteins of 70 and 66 kDa (Fig. 5B, lane 1). Two smaller bands have also been observed, in addition to the 90-kDa protein triplet, in whole-EB preparations (18), although these bands were larger than those reported here in COMCs. The reduction in size of these smaller proteins possibly resulted from proteolytic cleavage occurring during the preparation of the COMCs, an effect which could also account for the double MOMP band.

*pomp90A/B* 5'- atc gcc tgg ttg ggg ggg ggg ggg ... ggt acc tct cct gcc -3'  
 ile ala ser leu gly gly gly gly ... gly thr ser pro ala  
*pomp91A* 5'- gtc gcc tca ttg gga ggg ggg ggg gtc gcc cca gat cct gcc -3'  
 val ala ser leu gly gly gly gly val ala pro asp pro ala  
*pomp91B* 5'- atc gcc tgg ttg ggg ggg ggg ggg ... ggt acc tct cca gcc -3'  
 ile ala ser leu gly gly gly gly ... gly thr ala pro ala

FIG. 3. The polyguanine tract regions of the *pomp* genes. These sequences can be found under GenBank/EMBL accession no. U65942 for *pomp91A* (positions 1283 to 1324) and *pomp90A* (positions 3942 to 3980) and U65943 for *pomp91B* (positions 1828 to 1866) and *pomp90B* (positions 4481 to 4519). Dots represent gaps introduced into the sequences to maximize alignment.

Coomassie staining indicated that the proteins were present in low amounts in COMCs (Fig. 5A) and thus appear to be highly immunogenic and immunodominant: they were recognized by a large proportion of the postabortion sera obtained from individual ewes (for example, Fig. 5B, lanes 3 and 4). The bands were also recognized by the P90F31 and P91BF99 affinity-purified antibodies (Fig. 5B, lanes 5 and 6), confirming that these proteins are the products of the genes from which *p90f31* and *p91bf99* originated.

**Expression of the amino- and carboxy-terminal halves of POMP90A.** Immunoblotting of the expressed and purified products of both the 5' and 3' fragments of *pomp90A* showed that all 12 abortion and avian strain monoclonal antibodies and the pool of postabortion sera only detected the amino-terminal half (Fig. 6A and B). There was no reaction with the carboxy-terminal half, even when further postabortion sera were tested. In contrast, sera from sheep that were vaccinated with the purified carboxy-terminal product showed clear recognition of the 90-kDa protein triplet in immunoblotting of a whole-EB preparation (Fig. 6C).

As controls, the P90F31 and P91BF99 affinity-purified antibodies only recognized the amino-terminal product (data not shown), and a pool of negative control sera from a flock free of chlamydial infection (Fig. 6A, lanes 2 and 4) and anti-MOMP monoclonal antibody 4/11 (19) (data not shown) recognized neither product. Further analysis of the expressed products of *p90f31* and *p91bf99* with the avian monoclonal antibodies showed that three (including 89-76/049 in Fig. 6B, lane 1) recognized P90F31, while two (including 89-84/019 in Fig. 6B,




FIG. 4. Multiple sequence alignment of the predicted amino acid sequences of the *pomp* genes. Sequences were aligned by using the PILEUP program of the Wisconsin GCG package. Dots represent gaps introduced into the sequences to maximize sequence similarity. The blackened areas indicate amino acid identity, whereas the shaded residues indicate different levels of similarity (the darker shading represents a greater degree of similarity). P90F31 and P91BF99 are represented by residues 311 to 382 in POMP90A and POMP90B and residues 44 to 195 in POMP91B, respectively. Signal peptidase I cleavage sites were predicted for POMP90A, POMP90B, and POMP91B between positions 16 and 17 (arrowhead).

lane 3) recognized P91BF99 (data not shown). The remaining five avian strain (including 89-73/053 in Fig. 6B, lane 7) and three abortion strain (including 181 in Fig. 6B, lane 5) monoclonal antibodies must recognize the regions outside: the P90F31 and equivalent P91BF99 regions of the amino-terminal half of POMP90A. Although the size of the carboxy-terminal product, estimated by SDS-PAGE analysis, correlates well with its theoretical molecular mass of 48 kDa, the amino-terminal product ran anomalously, giving an estimated mass of 49 kDa, which is larger than its theoretical size of 41 kDa.

**Immunofluorescence.** Immunofluorescence microscopy of S26/3-infected McCoy cells at different times throughout the chlamydial growth cycle (every 6 h from 18 to 54 h plus 72 h postinfection) revealed weak immunofluorescent staining of very small inclusions at 18 h postinfection when anti-MOMP monoclonal antibody 4/11 was used. This staining increased in

intensity by 24 h, when inclusions were clearly recognizable within the cytoplasm of the McCoy cells, and by 30 h, inclusions were highly fluorescent. The inclusions remained highly fluorescent at all subsequent times up to 72 h postinfection. In contrast, with anti-POMP monoclonal antibodies 181 and 0040 (Table 1), weak immunofluorescent staining was not observed until 30 to 36 h postinfection and it was not until 42 h that inclusions were highly fluorescent. This staining intensity was maintained at all subsequent times up to 72 h postinfection. Figure 7 shows the immunofluorescent staining of inclusions at 48 h postinfection with both anti-POMP (A and B) and anti-MOMP (C) monoclonal antibodies (approximately one cell in three was infected). Nonspecific staining was not observed with any of these antibodies on uninoculated McCoy cells, and no staining was observed on infected cells with a nonspecific monoclonal antibody (Fig. 7D).




FIG. 5. Immunoblot analysis of EB outer membrane preparations. COMCs prepared as detailed in Materials and Methods were subjected to SDS-PAGE on 7.5% gels under reducing conditions and Coomassie stained (A) or immunoblotted (B) with the pool of postabortion sera used to screen the  $\lambda$ gt11 library from which *p90F31* and *p91Bf99* were isolated (lane 1); a pool of negative control sera from a flock free of chlamydial infection (lane 2); sera from two individual ewes, 3205 (lane 3) and 3210 (lane 4); and P91Bf99 (lane 5) and P90F31 (lane 6) affinity-purified antibodies. Molecular masses are indicated.

**Homologous genes in other chlamydial species.** Analysis of genomic DNAs from *C. trachomatis* (VW1 serovar D), *C. pneumoniae* (IOL-207), and four subtypes of *C. pecorum* (P787, W73, PV3056/3, and JP1751) by Southern blotting with *p91Bf99* and *pomp90A* probes under low-stringency hybridization and washing conditions failed to reveal any related sequences (Fig. 8, lane 4 for P787). With the FePn strain of *C. psittaci*, *Eco*RI fragments of 4.4 and ~20 kb hybridized weakly under low-stringency conditions with the *pomp90A* (Fig. 8, lane 3), but not the *p91Bf99* (data not shown), probe. However, with the avian strain of *C. psittaci*, *Eco*RI fragments of 2.5, 6.2, and ~14 kb showed strong homology with both the *pomp90A* (Fig. 8, lane 2) and *p91Bf99* (data not shown) probes, even under high-stringency conditions. No extra bands were identified by exposing blots to X-ray film for longer periods.

## DISCUSSION

The chlamydial outer membrane surface must play an important role in mucosal epithelial cell infection, being involved in the attachment of EBs to the host cell membrane, their subsequent uptake, and possibly the prevention of lysosome




FIG. 6. Immunoblot analysis of the expressed amino- and carboxy-terminal halves of POMP90A. (A and B) Purified amino (A, lanes 1 and 2; B, lanes 1, 3, 5, and 7) and carboxyl (A, lanes 3 and 4; B, lanes 2, 4, 6, and 8) fragments of POMP90A (0.3- and 0.2- $\mu$ g aliquots in A and B, respectively) were subjected to SDS-PAGE on 10% gels under reducing conditions and immunoblotted with the pool of postabortion sera (A, lanes 1 and 3); the pool of negative control sera (A, lanes 2 and 4); and anti-90-kDa monoclonal antibodies (B) 89-76/049 (lanes 1 and 2), 89-84/019 (lanes 3 and 4), 181 (lanes 5 and 6), and 89-73/053 (lanes 7 and 8). (C) Whole-EB preparations were subjected to SDS-PAGE on 7.5% gels under reducing conditions and immunoblotted with sera from ewes immunized with the carboxy-terminal half of POMP90A. Sera from ewes 2631 (lanes 1 and 2) and 2637 (lanes 3 and 4) collected preimmunization (lanes 1 and 3) and postimmunization (lanes 2 and 4) were used. Molecular masses are indicated.




FIG. 7. Immunofluorescence microscopy of *C. psittaci* S26/3-infected McCoy cells. *C. psittaci* S26/3-infected McCoy cells at 48 h postinfection were probed with putative outer membrane protein-specific monoclonal antibodies 181 (A) and 0040 (B) and MOMP-specific monoclonal antibody 4/11 (C) as detailed in Materials and Methods. A nonspecific (antileukotoxin) monoclonal antibody (D) was used as a negative control (leukotoxin is a cytolytic pore-forming protein from *Pasteurella haemolytica* serotype A1).

fusion with the inclusion membrane. The abundance and immunogenicity of the MOMP have caused most research to be focused on this protein. However, antibodies against other chlamydial proteins are produced during the course of a natural or experimental infection, particularly a group of three proteins with molecular masses of about 90 kDa (12, 18, 27). Additionally, a monoclonal antibody specifically recognizing an 89-kDa protein present in COMCs partially reduced infectivity in vitro (5). This 89-kDa protein band was reported later to consist of a triplet of proteins of 80, 85, and 90 kDa (27), which are the same as the 90-kDa protein triplet described in the present study from comparison of reactivities of antigen-




FIG. 8. Southern blot analysis of other *Chlamydia* subtypes and species. *Eco*RI-digested genomic DNAs from *C. psittaci* OEA (lane 1), avian (lane 2), and FePn (lane 3) strains and *C. pecorum* ovine arthritis strain P787 (lane 4) were Southern blotted and probed with DIG-labelled 1.1- and 1.6-kb fragments of *pomp90A* as described in Materials and Methods. Weak hybridization signals are indicated by arrowheads. DNA size markers (lane 5) are indicated.

specific monoclonal antibodies, affinity-purified antibodies, and a series of anti-90-kDa chlamydial parakeet strain monoclonal antibodies (32). These 90-kDa proteins are minor but highly immunogenic components of the COMC (Fig. 5) that we have previously shown to protect sheep from experimental chlamydial infection (30). They have also been suggested to be important for the diagnosis of OEA strain infection (27). For these reasons, it was considered extremely important to characterize these proteins at the molecular level since they may play an important role in both the serodiagnosis of, and protection from, OEA *C. psittaci*.

Previously, we have isolated two  $\lambda$ gt11 clones, containing *p90f31* and *p91Bf99*, which were shown to be part of a gene or genes coding for a group of three 90-kDa proteins (18). These bands could have resulted from separate related genes or from the posttranslational modification of one or more genes. Southern blot analysis of genomic DNA suggested several genes (18), a finding that was confirmed in this study, in which four genes have been identified and isolated. The genes were located as two pairs in different regions of the chlamydial genome (Fig. 2). Since the OEA *C. psittaci* genome has not been mapped, the genomic distance separating these two regions has not been determined, although it could be estimated by pulsed-field gel electrophoresis. The 3,202 bp of complete identity that exists between the two regions (Fig. 2) encompasses the *pomp90A* and *pomp90B* genes, as well as the 42 bp downstream and 640 bp upstream (includes the 130-bp intergenic region) DNA sequences (Fig. 2). This may indicate the point at which divergence in the sequence has occurred during evolution, resulting in the duplication of the genes. This duplication was shown to be genuine and not due to cloning artifacts caused by rearrangements occurring during the construction of the  $\lambda$ EMBL-3 library, by comparison of Southern hybridization restriction enzyme banding patterns of the lambda clones (Fig. 1) with those of genomic DNA (18), as well as by restriction endonuclease analysis of PCR-generated genomic DNA fragments containing the *pomp* gene sequences.

Sequence analysis predicts that the protein products of these *pomp* genes are located in the outer membrane. But of more importance are the immunofluorescence results obtained by using POMP-specific monoclonal antibodies (Fig. 7) and the immunoblotting results obtained with the Sarkosyl-DTT-insoluble COMC fraction (Fig. 5), since they provide experimental evidence of the localization of the 90-kDa proteins in the outer membrane. These results also imply a direct correlation between the surface accessibility of the proteins and *in vivo* immune recognition. Further evidence supporting the outer membrane location of these proteins came recently from Everett and Hatch, who investigated the probable cellular location of the cysteine-rich proteins (7). They clearly demonstrated the presence of the characteristic triplet of 90-kDa proteins in COMCs of *C. psittaci* 6BC (Fig. 1 in reference 7), as well as the surface labelling, albeit weak labelling, of these proteins using the hydrophobic affinity probe 3-(trifluoromethyl)-3-(*m*-[<sup>125</sup>I]iodophenyl)-diazirine (Fig. 4 in reference 7), although they did not discuss or refer to these results in their report.

Since the number of chlamydiae increases dramatically during infection, the inability to detect fluorescent staining at times earlier than 30 to 36 h postinfection with the anti-POMP monoclonal antibodies may be a consequence of the proteins being minor components of COMCs (i.e., due to detection levels). An alternative explanation could be that the proteins are products of late genes. Although the latter part of the cycle is asynchronous, with cells containing both dividing reticulate bodies and mature EBs, at 48 h chlamydiae are predominantly in the form of EBs, as judged by transmission electron micros-

copy. This implies that the fluorescence observed at this time (Fig. 7A and B), and at subsequent times postinfection, is due to staining of EBs and, possibly, reticulate bodies just prior to their reorganization into EBs. This requires further extensive investigation by electron microscopy at times early and late in the chlamydial cycle.

The cysteine content of the proteins ranges from 1.6 to 2% (POMP90A/B, POMP91A, and POMP91B contain 13, 15, and 17 cysteine residues, respectively), which is similar to the 1.9% of the MOMP and may indicate that the proteins are also involved in the disulfide-bonded cross-linking which is responsible for the structural rigidity of EBs, as was also suggested by Vretou et al. (33). Furthermore, the immunoblotting results obtained with the amino- and carboxy-terminal halves of POMP90A (Fig. 6), which show that the major immunological response was to the amino-terminal half of the protein, could indicate that the proteins form two domains with the carboxy-terminal domain inserted in the outer membrane and the amino-terminal domain exposed on the surface. Indeed, the short polyglycine region located centrally in each protein, coded for by the polyguanine tract (Fig. 3), may reflect a flexible linking region between two such protein domains. The lack of immune recognition of the carboxy-terminal fragment was investigated further by attempting to raise antibodies to the recombinant protein in sheep. Identification of the 90-kDa protein triplet by immunoblot analysis with sera from these immunized animals shows that the carboxyl-terminal fragment is, indeed, immunogenic, and so the lack of recognition with postabortion sheep sera is possibly due to the internal localization of this part of the protein. A further explanation could be that part of the carboxy-terminal end of the protein is surface exposed but inaccessible to antibody. The surface exposure of the amino-terminal fragment, suggested by immunofluorescence microscopy studies using monoclonal antibodies which recognize this part of the protein (Fig. 7), and the possible internal localization of the carboxy-terminal fragment require further investigation by immunoelectron microscopy using the anti-90-kDa monoclonal antibodies and monospecific antisera prepared against the carboxy-terminal protein fragment, respectively.

Since this group of proteins may play an important role in protective immunity against and/or the serodiagnosis of chlamydial infection, we searched for homologous genes in other chlamydial species (isolates of *C. pecorum*, *C. trachomatis*, and *C. pneumoniae*) but failed to detect any related gene sequences (Fig. 8). However, this does not necessarily imply that such genes are absent but that the homologies are too low for identification. The possible absence, or low homologies, of these genes in the *C. pecorum* subtypes should make the 90-kDa proteins suitable as serodiagnostic antigens, enabling ewes infected with the OEA agent to be identified without the problem of false-positive values due to widespread *C. pecorum* infection. The antigens are currently undergoing assessment for serodiagnostic potential, the preliminary results of which look promising (16). The role of the proteins in protection remains an open question, although they are highly immunogenic components of the subcellular vaccine that we have demonstrated to be protective (30).

This is the first report describing the isolation from *Chlamydia* species of duplicated genes coding for a cluster of immunogenic envelope proteins which may participate in key steps of the infection process. As research on these proteins is in its infancy and there are no homologous genes or proteins in the various databases, no function can be assigned. This is compounded by the constraints we have on the growth and genetic manipulation of the organism, which is perhaps the reason that the functions of other chlamydial proteins, includ-

ing the MOMP and Omp2, have yet to be definitively demonstrated. Regardless of this, the proteins are important serodiagnostic and vaccine antigen candidates that we believe will eventually have a broad impact on most chlamydial species.

#### ACKNOWLEDGMENTS

We thank J. Machell for providing EBs for Southern blot analysis, S. Wyllie for preparing the outer membrane EB fraction, I. E. Anderson for assistance in screening the  $\lambda$ EMBL-3 library, M. Livingstone for assistance with the sheep vaccinations, and B. J. Easter for photography. We are grateful to D. Buxton for helpful discussions. We also thank E. Vretou, Institut Pasteur Hellenique, Athens, Greece, for supplying the anti-POMP monoclonal antibodies; W. Donachie for the antileukotoxin antibody; and J. Trehanar, Institute of Ophthalmology, London, UK, for the *C. trachomatis* and *C. pneumoniae* isolates.

This work was supported by grants from the European Commission, Division of Agricultural Research, project CT93-0957 (AIR 3), and the Scottish Office Agriculture, Environment and Fisheries Department.

#### ADDENDUM

Since this report was submitted for publication, nine genes with protein sequences homologous to those of the POMP have been identified in *C. trachomatis* serovar D (D/UW-3/Cx) trachoma biovar by the Chlamydia Genome Project (on the World Wide Web at URL <http://chlamydia-www.berkeley.edu:4231/index.html>). The genes have been named *pmpA* to *pmpI*.

#### REFERENCES

- Aitken, I. D. 1991. Enzootic (chlamydial) abortion, p. 43-48. In W. B. Martin and I. D. Aitken (ed.), *Diseases of sheep*. Blackwell Scientific Publications, Oxford, England.
- Anderson, I. E., A. J. Herring, G. E. Jones, J. C. Low, and A. Greig. 1995. Development and evaluation of an indirect ELISA to detect antibodies to abortion strains of *Chlamydia psittaci* in sheep sera. *Vet. Microbiol.* 43:1-12.
- Baehr, W., Y.-X. Zhang, T. Joseph, H. Su, F. E. Nano, K. D. E. Everett, and H. D. Caldwell. 1988. Mapping antigenic domains expressed by *Chlamydia trachomatis* major outer membrane protein genes. *Proc. Natl. Acad. Sci. USA* 85:4000-4004.
- Caldwell, H. D., J. Kromhout, and J. Schachter. 1981. Purification and partial characterization of the major outer membrane protein of *Chlamydia trachomatis*. *Infect. Immun.* 31:1161-1176.
- Cevenini, R., M. Donati, E. Brocchi, F. De Simone, and M. La Placa. 1991. Partial characterization of an 89-kDa highly immunoreactive protein from *Chlamydia psittaci* A/22 causing ovine abortion. *FEMS Microbiol. Lett.* 81: 111-116.
- Collett, B. A., W. J. Newhall V, R. A. Jersild, Jr., and R. B. Jones. 1989. Detection of surface-exposed epitopes on *Chlamydia trachomatis* by immune electron microscopy. *J. Gen. Microbiol.* 135:85-94.
- Everett, K. D. E., and T. P. Hatch. 1995. Architecture of the cell envelope of *Chlamydia psittaci* 6BC. *J. Bacteriol.* 177:877-882.
- Fleischmann, R. D., M. D. Adams, O. White, R. A. Clayton, E. F. Kirkness, A. R. Kerlavage, C. J. Bult, J.-F. Tomb, B. A. Dougherty, J. M. Merrick, K. McKenney, G. Sutton, W. FitzHugh, C. Fields, J. D. Gocayne, et al. 1995. Whole-genome random sequencing and assembly of *Haemophilus influenzae* Rd. *Science* 269:496-512.
- Fox, A., J. C. Rogers, J. Gilbert, S. Morgan, C. H. Davis, S. Knight, and P. B. Wyrick. 1990. Muramic acid is not detectable in *Chlamydia psittaci* or *Chlamydia trachomatis* by gas chromatography-mass spectrometry. *Infect. Immun.* 58:835-837.
- Fraser, C. M., J. D. Gocayne, O. White, M. D. Adams, R. A. Clayton, R. D. Fleischmann, C. J. Bult, A. R. Kerlavage, G. Sutton, J. M. Kelley, J. L. Fritchman, J. F. Weidman, K. V. Small, M. Sandusky, J. Fuhrmann, et al. 1995. The minimal gene complement of *Mycoplasma genitalium*. *Science* 270:397-403.
- Garnier, J., D. J. Osguthorpe, and B. Robson. 1978. Analysis of the accuracy and implications of a simple method for predicting the secondary structure of globular proteins. *J. Mol. Biol.* 120:97-120.
- Griffiths, P. C., H. L. Philips, M. Dawson, and M. J. Clarkson. 1992. Antigenic and morphological differentiation of placental and intestinal isolates of *Chlamydia psittaci* of ovine origin. *Vet. Microbiol.* 30:165-177.
- Herring, A. J., I. E. Anderson, M. McClenaghan, N. F. Inglis, H. Williams, B. A. Matheson, C. P. West, M. Rodger, and R. P. Brett. 1987. Restriction endonuclease analysis of DNA from two isolates of *Chlamydia psittaci* obtained from human abortions. *Br. Med. J.* 295:1239.
- Herring, A. J., T. W. Tan, S. Baxter, N. F. Inglis, and S. Dunbar. 1989. Sequence analysis of the major outer membrane protein gene of an ovine abortion strain of *Chlamydia psittaci*. *FEMS Microbiol. Lett.* 65:153-158.
- Herring, A. J., M. C. McCafferty, G. E. Jones, S. Dunbar, and A. A. Andersen. 1994. Vaccination against chlamydial abortion in sheep: problems and progress with a recombinant vaccine, p. 118-121. In J. Orfila, G. I. Byrne, M. A. Chernesky, J. T. Grayston, R. B. Jones, G. L. Ridgway, P. Saikku, J. Schachter, W. E. Stamm, and R. S. Stephens (ed.), *Chlamydial infections*. Societa Editrice Esculapio, Bologna, Italy.
- Jones, G. E., J. Machell, E. Vretou, and D. Longbottom. Unpublished data.
- Klein, P., M. Kanehisa, and C. DeLisi. 1985. The detection and classification of membrane spanning proteins. *Biochim. Biophys. Acta* 815:468-476.
- Longbottom, D., M. Russell, G. E. Jones, F. A. Lainson, and A. J. Herring. 1996. Identification of a multigene family coding for the 90 kDa proteins of the ovine abortion subtype of *Chlamydia psittaci*. *FEMS Microbiol. Lett.* 142:277-281.
- McCafferty, M. C., A. J. Herring, A. A. Andersen, and G. E. Jones. 1995. Electrophoretic analysis of the major outer membrane protein of *Chlamydia psittaci* reveals multimers which are recognized by protective monoclonal antibodies. *Infect. Immun.* 63:2387-2389.
- McClenaghan, M., A. J. Herring, and I. D. Aitken. 1984. Comparison of *Chlamydia psittaci* isolates by DNA restriction endonuclease analysis. *Infect. Immun.* 45:384-389.
- McClenaghan, M., N. F. Inglis, and A. J. Herring. 1991. Comparison of isolates of *Chlamydia psittaci* of ovine, avian and feline origin by analysis of polypeptide profiles from purified elementary bodies. *Vet. Microbiol.* 26: 269-278.
- McGeoch, D. J. 1985. On the predictive recognition of signal peptide sequences. *Virus Res.* 3:271-286.
- Nakai, K., and M. Kanehisa. 1991. Expert system for predicting protein localization sites in gram-negative bacteria. *Proteins* 11:95-110.
- Novagen Inc. 1995. pET system manual, 6th ed. Novagen Inc., Madison, Wis.
- Raulston, J. E. 1995. Chlamydial envelope components and pathogen-host cell interactions. *Mol. Microbiol.* 15:607-616.
- Salari, S. H., and M. E. Ward. 1981. Polypeptide composition of *Chlamydia trachomatis*. *J. Gen. Microbiol.* 123:197-207.
- Souriau, A., J. Salinas, C. De Sa, K. Layachi, and A. Rodolakis. 1994. Identification of subspecies- and serotype 1-specific epitopes on the 80- to 90-kilodalton protein region of *Chlamydia psittaci* that may be useful for diagnosis of chlamydial induced abortion. *Am. J. Vet. Res.* 55:510-514.
- Struyve, M., M. Moons, and J. Tommassen. 1991. Carboxy-terminal phenylalanine is essential for the correct assembly of a bacterial outer membrane protein. *J. Mol. Biol.* 218:141-148.
- Su, H., Y.-X. Zhang, O. Barrera, N. G. Watkins, and H. D. Caldwell. 1988. Differential effect of trypsin on infectivity of *Chlamydia trachomatis*: loss of infectivity requires cleavage of major outer membrane protein variable domains II and IV. *Infect. Immun.* 56:2094-2100.
- Tan, T.-W., A. J. Herring, I. E. Anderson, and G. E. Jones. 1990. Protection of sheep against *Chlamydia psittaci* infection with a subcellular vaccine containing the major outer membrane protein. *Infect. Immun.* 58:3101-3108.
- von Heijne, G. 1986. A new method for predicting signal sequence cleavage sites. *Nucleic Acids Res.* 14:4683-4690.
- Vretou, E., H. Loutrari, L. Mariani, K. Costelidou, P. Eliades, G. Conidou, S. Karamanou, O. Mangana, V. Siarkou, and O. Papadopoulos. 1996. Diversity among abortion strains of *Chlamydia psittaci* demonstrated by inclusion morphology, polypeptide profiles and monoclonal antibodies. *Vet. Microbiol.* 51:275-289.
- Vretou, E., P. Yannikopoulou, E. Psarron, P. Papavassiliou, V. Pallini, and L. Bini. 1996. 2-D electrophoretic and antigenic characterisation of the protein family at 90 kDa in the envelope of *C. psittaci* serotype 1, p. 52. In A. Stary (ed.), *Proceedings to the Third Meeting of the European Society for Chlamydia Research*. Societa Editrice Esculapio, Bologna, Italy.